



Faculty of Resource Science and Technology

**DETERMINATION OF GENETIC RELATEDNESS IN KELAMPAYAN
PLANTLETS REGENERATED FROM TISSUE CULTURE USING DAMD
MARKERS**

Suthashinikisan A/P Krishnan

**Bachelor of Science with Honours
(Biotechnology Resource)
2013**

**DETERMINATION OF GENETIC RELATEDNESS IN KELAMPAYAN PLANTLETS
REGENERATED FROM TISSUE CULTURE USING DAMD MARKERS**

Suthashinikisan A/P Krishnan

**This thesis is submitted in partial fulfillment of the requirements for the degree of
Bachelor of Science with Honours
(Resource Biotechnology)**

**Faculty of Resource Science and Technology
UNIVERSITY MALAYSIA SARAWAK
2013**

ACKNOWLEDGEMENTS

Above all, I would like to thank God for His blessings and guidance in the completion of this project. I would like to express my special thanks of gratitude to my supervisor, Dr. Ho Wei Seng and co-supervisor Dr. Pang Shek Ling for giving me opportunity to do this project to complete my Final Year project and for their guidance throughout in the completion of this research.

I would also like to thank Ms Kamalia, Ms Grace Ting Jen Ching (Msc Student) and Ms. Natalie Gali (Msc Student) for their kindness.

Last but not least, I would also like to thank my parents and friends who helped me a lot in finishing this project within the limited time.

DECLARATION

I hereby declare that this is based on original work except for quotations and citations which have been acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UNIMAS or other institutions.

.....

SUTHASHINIKISAN KRISHNAN, 28406

Date:

Resource Biotechnology

Department of Molecular Biology

Faculty of Resource Science and Biotechnology

Universiti Malaysia Sarawak (UNIMAS)

TABLE OF CONTENTS

	Page
ACKNOWLEDGEMENTS	I
DECLARATION	II
TABLE OF CONTENTS	III
LIST OF ABBREVIATIONS	V
LIST OF FIGURES	VI
LIST OF TABLES	IX
ABSTRACT	X
CHAPTER I INTRODUCTION	1
CHAPTER II LITERATURE REVIEW	4
2.1 <i>Neolamarckia cadamba</i>	4
2.2 <i>Neolamarckia macrophylla</i>	5
2.3 Direct Amplification of Minisatellite-region DNA	7
2.4 Tissue Culture Variations	8
CHAPTER III MATERIALS AND METHODS	10
3.1 Plant samples and Isolated DNA Samples	10
3.2 Polymerase Chain Reaction Amplification	11
3.2.1 PCR Condition	11
3.3 Agarose Gel Electrophoresis	13
3.4 DNA Scoring and Binary Data Analysis	13
CHAPTER IV RESULTS	15
4.1 Genomic DNA Screening and Quantification	15
4.2 PCR Optimization and Amplification	17
4.2.1 PCR Optimization of <i>N. cadamba</i>	17
4.2.2 PCR Optimization of <i>N. macrophylla</i>	20
4.2.3 <i>Neolamarckia cadamba</i> DNA Amplification with M13, 33.6 and YNZ-22 Primers	23
4.2.4 <i>Neolamarckia macrophylla</i> DNA Amplification with M13, 33.6 and YNZ Primers	25
4.3 Genetic Relatedness	27
4.3.1 Genetic Variation among <i>Neolamarckia</i>	27

	<i>cadamba</i> Tissue Cultured samples	
4.3.1.1	M13 universal primer	27
4.3.1.1	33.6 universal primer	28
4.3.1.1	YNZ-22 universal primer	30
4.3.1.1	Based on all three M13, 33.6 and YNZ-22 primers	31
4.3.2	Genetic Variation among <i>Neolamarckia</i> <i>macrophylla</i> Tissue Cultured samples	33
4.3.1.1	M13 universal primer	33
4.3.1.1	33.6 universal primer	35
4.3.1.1	YNZ-22 universal primer	36
4.3.1.1	Based on all three M13, 33.6 and YNZ-22 primers	37
CHAPTER V	DISCUSSIONS	39
5.1	Genomic DNA Screening and Quantification	39
5.2	PCR Optimization and Amplification	39
5.3	Genetic Diversity among Tissue Cultured	41
5.3.1	Genetic Relatedness among <i>N. cadamba</i> Tissue Cultured samples	41
5.3.2	Genetic Relatedness among <i>N. macrophylla</i> Tissue Cultured samples	42
CHAPTER VI	CONCLUSIONS AND RECOMMENDATIONS	43
REFERENCES		44
APPENDICES		47

LIST OF ABBREVIATIONS

bp	basepair
cm	centimeter
CTAB	Cetyl Trimethyl Ammonium Bromide
DNA	deoxyribonucleic acid
dNTPs	deoxynucleotide-triphosphates
kb	kilo basepair
m	meter
MgCl₂	Magnesium chloride
min	minutes
ml	milliliter
mm	millimeter
mM	milimolar
μl	microliter
PCR	Polymerase Chain Reaction
pH	potential of Hydrogen
pmol/μl	picomolar per microliter
POPGENE	Population genetics
TAE	Tris- Acetate –EDTA
U	Unit
V	Volt
%	percent
°C	Celcius

LIST OF FIGURES

Figure	Page
2.1 Flowers of <i>N. cadamba</i> .	4
2.1 Flowers of <i>N. macrophylla</i> .	4
2.3 Seven-month old <i>N. macrophylla</i> leaf.	6
3.1 An example of selected bands generated by M13 universal primer named after the primer and a hyphenated number corresponding to the order of their migrations.	14
3.2 An example of selected bands generated by M13 universal primer named after the primer and a hyphenated number corresponding to the order of their migrations based on gel photo produced using AlphaEaseFC Software.	14
4.1 DNA screening of <i>N. cadamba</i> DNA samples. Total of six samples from the tissue cultured sample are represented in the photograph. Lambda <i>Hind</i> III DNA marker was used. Lane 1 to 6 represents MGK03, MGK04, MGK05, MGK06, MGK08 and MGK10	15
4.2 DNA screening of <i>N. macrophylla</i> DNA samples. Total of nine samples from the tissue cultured sample are represented in the photograph. Lambda <i>Hind</i> III DNA marker was used. Lane 1 to 9 represents sample MGR2-6, MGR6-4, MGR10-2, MGR10-6, MGR6-3A, MGR10-1, MGR2-1, MGR2-2 and MGR6-3B	16
4.3 PCR amplification with different template DNA concentrations. Sample MGK03 of <i>N. cadamba</i> was used for the optimization. (a) Lane 1 to 10 represents DNA concentration of 5ng, 10ng, 15ng, 30ng, 45ng, 50ng, and 100ng. (b) Lane 1 to 6 represents DNA concentration of 1ng with different <i>Taq</i> polymerase and primer volume.	18
4.4 PCR amplification with different temperature profile. Sample MGR6-3A of <i>N. macrophylla</i> was used for the optimization. (a) Optimization using 33.6 primer. (b) Optimization using YNZ-22 primer. Lane 1 to 8 represents temperature ranging at 46.1 °C, 46.7 °C, 47.2 °C, 47.6 °C, 48.0 °C, 48.7 °C, 49.4 °C, and 50.1 °C	21
4.5 DAMD-PCR profiles of <i>N. cadamba</i> using M13 universal primer. A 100 bp	23

- and 1 kb DNA ladder was used as a DNA molecular weight marker. Lane 1 to 6 represents MGK03, MGK04, MGK05, MGK06, MGK08 and MGK10 follows in order
- 4.6 DAMD-PCR profiles of *N. cadamba* using 33.6 primer (Lane 1 to 6 24 represents MGK03, MGK04, MGK05, MGK06, MGK08 and MGK10 follows in order) and YNZ-22 primer (Lane A to F MGK03, MGK04, MGK05, MGK06, MGK08 and MGK10 follows in order). A 1 kb ladder was used as a DNA molecular weight marker
- 4.7 DAMD-PCR profiles of *N. macrophylla* using M13 universal primer. A 100 24 bp ladder was used as a DNA molecular weight marker. Lane 1 to 9 represents sample MGR2-6, MGR6-4, MGR10-2, MGR10-6, MGR6-3A, MGR10-1, MGR2-1, MGR2-2 and MGR6-3B
- 4.8 DAMD-PCR profiles of *N. macrophylla* using 33.6 primer. A 100 bp ladder 25 was used as a DNA molecular weight marker. Lane 1 to 9 represents sample MGR2-6, MGR6-4, MGR10-2, MGR10-6, MGR6-3A, MGR10-1, MGR2-1, MGR2-2 and MGR6-3B
- 4.9 DAMD-PCR profiles of *N. macrophylla* using YNZ-22 primer. A 1 kb ladder 26 was used as a DNA molecular weight marker. Lane 1 to 9 represents sample MGR2-6, MGR6-4, MGR10-2, MGR10-6, MGR6-3A, MGR10-1, MGR2-1, MGR2-2 and MGR6-3B
- 4.10 UPGMA cluster analysis of DAMD similarities of 6 tissue culture samples of 28 *Neolamarckia cadamba* using M13 primer
- 4.11 UPGMA cluster analysis of DAMD similarities of 6 tissue culture samples of 29 *Neolamarckia cadamba* using 33.6 primer
- 4.12 UPGMA cluster analysis of DAMD similarities 6 tissue culture samples of 31 *Neolamarckia cadamba* using YNZ-22 primer
- 4.13 UPGMA cluster analysis of similarities among 6 tissue culture samples of *N. 32 cadamba*. The dendograms was analyzed by using M13, 33.6 and YNZ-22 minisatellite primers
- 4.14 UPGMA cluster analysis of DAMD similarities of 9 tissue culture samples of 34 *N. macrophylla* using M13 primer
- 4.15 UPGMA cluster analysis of DAMD similarities of 9 tissue culture samples of 36 *N. macrophylla* using 33.6 primer

- 4.16 UPGMA cluster analysis of DAMD similarities of 9 tissue culture samples of *N. macrophylla* using YNZ-22 primer 37
- 4.17 Cluster analysis of similarities among 6 tissue culture samples of *N. macrophylla*. The dendograms was analyzed by using M13, 33.6 and YNZ-22 primers 38

LIST OF TABLES

Table	Page
2.1 Morphological characteristics differences of <i>N. macrophylla</i> and <i>N. cadamba</i>	6
2.2 List of primers used in this study	8
3.1 List of samples of <i>N. cadamba</i> and <i>N. macrophylla</i>	10
3.2 PCR ingredients for 25 µl reaction mixture	11
3.3 A thermal cycling profile used for preliminary PCR optimization step	12
3.4 The PCR reaction conditions for PCR amplifications with M13 minisatellite universal primer.	12
3.5 The PCR reaction conditions for PCR amplifications with 33.6 and YNZ-22 minisatellite primer.	13
4.1 The optimum PCR reaction condition for PCR amplification with M13 universal primer on <i>N. cadamba</i>	19
4.2 The optimum PCR reaction condition for PCR amplification with 33.6 and YNZ primers	19
4.3 The optimum PCR reaction condition for PCR amplification with M13, 33.6 and YNZ-22 primers on <i>N. macrophylla</i>	20
4.4 Optimal thermal cycling profiles for both PCR amplification with 33.6 and YNZ-22 primers	22
4.5 Summary of genetic variation statistics for all M13 primer loci	27
4.6 Summary of genetic variation statistics for all 33.6 primer loci	29
4.7 Summary of genetic variation statistics for all YNZ-22 primer loci	30
4.8 Summary of genetic variation statistics for all loci	31
4.9 Summary of genetic variation statistics for all M13 primer loci	33
4.10 Summary of genetic variation statistics for all 33.6 primer loci	35
4.11 Summary of genetic variation statistics for all YNZ-22 primer loci	36
4.12 Summary of genetic variation statistics for all primer loci	38

Determination of Genetic Relatedness in Kelampayan Plantlets Regenerated from Tissue Culture using DAMD Markers

Suthashinikisan A/P Krishnan

Resource Biotechnology
Faculty of Resource Science and Technology
University Malaysia Sarawak

ABSTRACT

Neolamarckia cadamba and *Neolamarckia macrophylla* belongs to Rubiaceae family and locally known as white and red kelampayan respectively. They are economically important woody trees. These trees are large, deciduous and fast growing tree species. Most important of any *in vitro* propagation system is mass multiplication of plantlets that are genetically homogenous and phenotypically uniform. Modified CTAB method was used to extract the DNA. Directed amplification of Minisatellite-region DNA primers namely M13, YNZ, and 33.6 primers was used as a tool to assess the genetic relatedness of *in vitro* propagated *N. cadamba* and *N. macrophylla* hardened explants. The presence or absence of the amplification product showed allelic variation among individuals after visualizing as a band during electrophoresis. Cluster analysis of DAMD-PCR data using UPGMA (unweighted pair group method with arithmetic average) revealed the genetic relatedness among the species. A total of 21 bands and 24 bands were produced respectively. Mean gene diversity of *N. cadamba* and *N. macrophylla* is 0.1825 and 0.0926 respectively. Shannon index is 0.2632 and 0.1324 respectively. The low figure of both mean of gene diversity and Shannon index obtained shows that *N. cadamba* are related to each other while *N. macrophylla* samples are closely related. The UPGMA cluster analyses revealed *N. macrophylla* samples are highly related among the samples but two samples; MGR10-6 and MGR2-2 belongs to different clones shows close relatedness. The close relatedness among the samples indicates that tissue culture generated samples of both the kelampayan species shows very little polymorphism and therefore *in-vitro* regenerated plantlets in this study maintain their genetic integrity.

Keywords: Kelampayan, Random Amplified Polymerase DNA, *in vitro* propagation, genetic relatedness.

ABSTRAK

Neolamarckia cadamba dan *Neolamarckia macrophylla* tergolong dalam keluarga Rubiaceae dan nama tempatan masing-masing dikenali sebagai kelampayan putih dan merah. Pokok-pokok ini amat penting dari segi ekonomi. Hal ini kerana pokok ini besar, berdaun lebat serta tumbesaran yang cepat. Perkara penting dalam sistem pembiakan vitro adalah anak pokok yang seragam secara genomik dan fenotip meningkat dengan banyak. Kaedah CTAB telah digunakan untuk mengekstrak DNA. Primer M13, YNZ, dan 33.6 telah digunakan untuk menilai pertalian genetik diantara *N. cadamba* dan *N. macrophylla*. Analisis kelompok data DAMD menggunakan UPGMA telah menunjukkan pertalian genetik antara spesies. Min persamaan genetic bagi kumpulan pokok *N. cadamba* dan *N. macrophylla* ialah 0,1825 dan 0,0926. Manakala indeks Shannon adalah 0.2632 dan 0.1324. Analisis kelompok UPGMA di antara anak pokok tersebut menunjukkan bahawa kesemua anak pokok kedua-dua spesies tersebut menunjukkan hubungan rapat dari segi genomic. Namun dua sampel dari *N. macrophylla* (MGR10-6 dan MGR2-2) menunjukkan hubungan rapat walaupun diperolehi dari klon yang berbeza. Pertalian erat di kalangan sampel menunjukkan kedua-dua spesies kelampayan menunjukkan 'polymorphism' yang rendah dan oleh itu, tanaman kaedah *in vitro* yang digunakan dalam kajian ini mengekalkan integriti genetik mereka.

Kata kunci: Kelampayan, Hubungan genetic, Random Amplified Polymerase DNA, Kaedah *in vitro*.

CHAPTER I

INTRODUCTION

This research focus on the determination of the genetic relatedness of kelampayan plantlets: *Neolamarckia cadamba* and *Neolamarckia macrophylla*. Both this species belongs to Rubiaceae family, is an economically important species. It is commonly known as kelampayan, important for pulp and paper industry.

It is widely distributed from India, Nepal, through Thailand and Indo-China and eastward in the Malaysian Archipelago to Papua New Guinea (Joker, 2000). The tree is mostly found below 1000 m altitude and places where more than 1500 mm rain/ year are noted and also it can grow in dry areas with as little as 200 mm rain/year within the area of its natural distribution (Joker, 2000).

There are several studies used to assess the genetic relatedness of *in vitro* derived clones namely by karyological analysis, isozyme markers and polymerase chain reaction in conjunction with short primers. But most strategies have its limitations. For example, karyological analysis cannot reveal alterations in specific genes or small chromosomal rearrangements (Isabel *et al.*, 1993). Though isozyme markers provide a convenient method for detecting genetic changes, but are subjected to ontogenic variations. Also, only DNA regions coding for soluble proteins can be sampled.

PCR with short primers of arbitrary sequence (Williams *et al.*, 1990) shown to be sensitive for detecting variations among individuals between and within species (Carlson *et al.*,

1991; Roy *et al.*, 1992). This technique requires no prior knowledge of the genome and it needs only a small amount of DNA. Using this technique polymorphism can be detected in closely related organism.

Micropropagation of tree species offers rapid means of producing clonal planting stock for afforestation, woody biomass, and conservation of elite germplasm. Also, it is an alternative tool for crop improvement (Vasil, 1988). Therefore, tissue culture plays important role in replanting forest trees such as *N. cadamba* and *N. macrophylla* (Ndoye *et al.*, 2003).

Plantlets derived from *in vitro* culture might exhibit somaclonal variation (Larkin and Scowcroft, 1981) which is often heritable (Breiman *et al.*, 1987). The variations might be due to genotypes used (Breiman *et al.*, 1987), pathways of regeneration, and parameters used during *in vitro* culture such as morphology and cytology (Swedlund and Vasil, 1985) and molecular studies (Breiman *et al.*, 1989; Chawdhury *et al.*, 1994; Shenoy and Vasil, 1992).

Kelampayan improvement through traditional method is a slow, time and labor consuming process. So the optimization regeneration system is an important process and is considered as a prerequisite step towards plant transformation. Cell and tissue culture related to variability and selection efficiency are two essential components of the molecular breeding program (Lichtenstein and Draper, 1985).

The genetic analysis of these species derived from tissue culture is important because it will enable to choose the best quality trees with superior growth potential. Higher multiplication rates of selected tree genotypes are possible with micropropagation with uniform genotypes, resulting in short-term production gains. *N. cadamba* has been micropropagated because of its importance in the pulp and mining industries. Knowledge on the genetic stability of the proposed

samples will enable to plant more trees that show high good quality trees. Good quality includes larger width, high and faster growth.

However, the pressure for its valuable timber and fodder due to human activities has increased lately. Many tree felled off due to its importance in pulp and paper industry. Therefore, importance of replanting this tree has increased lately. Total of 18,851 ha of kelampayan trees planted in Sarawak which are approximately 8 % of overall species planted (Forest Department of Sarawak, 2012).

Today demand for plants with superior growth potential is high. Thus application of clonal propagation is widely used to produce high multiplication rates of uniform genotypes. Therefore, the objective of this study is to assess the genetic relatedness of *N. cadamba* and *N. macrophylla* plantlets derived from tissue culture with DAMD markers and its ability to form a uniform genetic profile. Also, to provide knowledge on genetic variance among the tested samples.

CHAPTER II

LITERATURE REVIEW

2.1 *Neolamarckia cadamba*

Neolamarckia cadamba or locally known as kelampayan is a timber species belongs to Rubiaceae family. The tree can grow up to 45 m tall and with diameter of 100 – 160 cm (Joker, 2000). The crown is open and round and bears drooping branches. The stem is straight and more or less cylindrical. Kelampayan has its flowers in orange color, small in dense and with globose heads while its leaves are up to 32 cm long (Joker, 2000).

N. cadamba (Roxb.) Bosser is a multipurpose tree known for many uses includes paper, wooden shoes, match sticks, plywood, pencil slats, wood carving, packing cases and boxes, and also splint manufacture (Lugo and Figueroa, 1984). The flowers of *N. cadamba* are offered at hindu shrines, and the fruits are eaten in India (Lugo and Figueroa, 1984).



Figure 2.1: Flowers of *N. cadamba*



Figure 2.2: Green leaves of *N. cadamba*

This tree is a fast growing tree and suitable for reforestation conservation especially in watersheds and also for windbreaks in agroforestry systems (Joker, 2000). *N. cadamba* is known for its lightweight hardwood which is poor durability therefore mainly used for pulp (producing low and medium quality paper) and veneer plywood. In addition, the bark and leaves have various medicinal uses such as astringent anti-hepatotoxic, antidiuretic, wound healing, antiseptic and anthelmintic (Patel and Kumar, 2008).

2.2 *Neolamarckia macrophylla*

Neolamarckia macrophylla (Roxb.) Bosser is a member of Rubiaceae family or more commonly known as red kelampayan. It has reddish green hairy leaf surface and its stem is in dark red. Red kelampayan is one type of fast-growing plants with a stem circumference more than 150 cm (diameter greater than 50 cm) (Halawane *et al.*, 2012). *N. macrophylla* is a tree with straight trunk that is free from branch up to 70% -80%. It drops the bottom leaves to grow straight up without branching and has smooth texture and straight wood grain. This tree grows well in locations with a height of 10-1000 m above sea level. Ability to grow on degraded land is also quite good, and even can be used as a buffer zone for the conservation purpose due to its deep roots (Halawane *et al.*, 2012).

With the red color and a smooth texture, this wood can be used as a raw material for plywood, furniture, plywood (veneer), home accessories and more. Also used as the raw material of short fiber pulp with medium quality.



Figure 2.3: Seven-month old *N. macrophylla* leaf

(Source: <http://jabon.web.id/jawaban-atas-pertanyaan-dari-jabon-merah/>)

Table 2.1: Morphological characteristics differences of *N. macrophylla* and *N. cadamba*

Characteristics	<i>N. macrophylla</i>	<i>N. cadamba</i>
Young leaf buds	Red	Brown
Basal leaves	Sharp-pointed	Flat
Primary leaf veins	Red	Yellowish green
Young stems	Blackish red	Brownish green
Mature tree trunks	Black	Brown grey
Fruit color	Ripen fruit is reddish brown	Ripen fruit is yellow

Source: Halawane *et al.*, 2012.

2.3 Directed Amplification of Minisatellite-region DNA (DAMD) Marker

Among the PCR based DNA marker systems, RAPD, DAMD and ISSR are commonly and extensively used tools for assessment of variability in crops. Due to their ease, rapidity and reliability, this marker system is efficient for analysis of molecular differentiation and for resolving taxonomic problems in plants (Ranade et al., 2006). Repetitive DNA sequences such as microsatellites, minisatellites, and satellites can be of great value in assessing a high level of polymorphism as they distributed throughout the genomes analyzed thus far.

Directed amplification of minisatellite-region DNA (DAMD) uses VNTR core sequences such as M13 as primers in PCR reactions (Heath et al. 1993). Minisatellites are tandemly repeated DNA regions of eukaryotic genomes. It shows high levels of variation due to differences in the number of repeated units (Jeffreys *et al.* 1985). It involved an inversion of a portion of a minisatellite DNA array with a single primer makes PCR is possible for the amplification of minisatellite core region (Heath *et al.* 1993). Since minisatellite core sequences which are used as primers are longer than RAPD-PCR primers, DAMD-PCR can be effectively carried out at relatively high stringency reactions (Bebeli *et al.* 1997).

However, this marker does have its limitation in terms of uncertain homology of co-migrating fragments in gel electrophoresis and reproducibility. This can be minimized by carefully adjusting the reaction and detection conditions. Table 2.2 shows the primers used in this study.

Table 2.2: List of primers used in this study

Primer code	Primer Sequence 5' to 3'	Nucleotide length
M13	5'- TTATGAAACGACGGCCAGT-3	19
33.6	5'- AGGGCTGGAGG - 3'	11
YNZ -22	5'- CTCTGGGTGTGGTGC- 3'	15

2.4 Tissue culture induced variations

Plant tissue culture has become one of the fundamental tools of plant science research. It is extensively employed in the production, conservation and improvement of plant resources. Therefore, *in vitro* culture techniques provide an alternative means of plant propagation and a tool for crop improvement (Vasil, 1988).

Tissue culture and *in vitro* plantlet regeneration systems provide an alternative production system for mass propagation of the desired plant genotype for commercial planting. These systems are applied when the selected genotype is required at mass scale, for commercial cultivation when the natural propagation does not meet the required demands.

However, *in vitro* techniques do induce somaclonal variations. According to Rathore *et al.*, (2011), the frequency of somaclonal variations varies with the source of explants and their regeneration pattern, media composition, and cultural conditions. These variations remain one of the major problems of many tissue-cultured plants (Rathore *et al.*, 2011). Uncontrolled and random spontaneous variations during culture process are major causes.

Nevertheless, variations are not always bad, it does sought for crop improvement by creation of novel variants through applications of advanced biotechnological tools (Rathore *et*

al., 2011). In addition, it induces variations in genetic manipulation of crops with phylogenetic traits. Since the somaclonal variations in tissue culture plantlets are possible, therefore the need to test and verify the genetic stability of regenerated plantlets is important. The variations can be determined at the morphological, cytological, biochemical, and molecular levels. At present, molecular markers are used in the analysis of genetic fidelity of *in vitro* propagated plantlets.

The most important part of any *in vitro* propagation system is mass multiplication of plantlets that are genetically homogenous and phenotypically uniform. Several approaches have been applied for identifying variants among micropropagated plants. Molecular markers have been shown to enhance breeding efforts in annual and perennial crops, since they are not altered by major environmental factors.

CHAPTER III

MATERIALS AND METHODS

3.1 Plant Samples and Isolated DNA Samples

Hardened tissue cultured kelampayan leaves of *N. cadamba* and *N. macrophylla* were obtained from the MGene Propagation Sdn Bhd. There are total of six samples of *N. cadamba* and nine samples of *N. macrophylla*. Table 3.1 shows the list of samples.

Table 3.1: List of samples of *N. cadamba* and *N. macrophylla*

			Species					
<i>Neolamarckia macrophylla</i>			<i>Neolamarckia cadamba</i>					
Clone 1	Clone 2	Clone 3	Clone 1	Clone 2	Clone 3	Clone 4	Clone 5	Clone 6
MGR	MGR	MGR	MGK	MGK	MGK	MGK	MGK	MGK
2-1	6-3A	10-1	03	04	05	06	08	10
MGR	MGR	MGR						
2-2	6-3B	10-2						
MGR	MGR	MGR						
2-6	6-4	10-6						

Total genomic DNA of *N. cadamba* and *N. macrophylla* was extracted by following modified CTAB method by Doyle and Doyle (1990). Purification of DNA sample is done by following protocol of Wizard® Genomic DNA Purification Kit (Promega). The presence of isolated genomic DNA from the leaf was determined by running a 0.8% agarose gel. The yield of DNA was measured using Nanodrop 2000 Spectrophotometer (Thermo Fisher Scientific, USA). The purity of DNA was determined by calculating the ratio of absorbance at 260 nm to that of 280 nm.

3.2 PCR Amplification

3.2.1 PCR Condition

Prior to begin the PCR reaction, a total of 25 µl PCR mixtures were prepared. PCR amplifications were performed on a Gradient Palm-CyclerTM (Corbett Research, Australia). The PCR products were checked using 1.5% agarose gel and run simultaneously with 100 bp or 1 kb DNA ladder (Promega, USA). The PCR optimization was carried out to determine the optimal condition for this analysis. The PCR reaction mixture profile in a total volume of 25 ul is shown in Table 3.1.

Table 3.2: PCR ingredients for 25 µl reaction mixture

Reagent	Concentration
Double distilled Water	-
PCR buffer	10 ×
dNTPs	2 mM
MgCl ₂	50 mM
Primer	2.5 pmol
<i>Taq</i> DNA polymerase	0.5 U
Template DNA	15 ng/ µl

The thermal cycling profile used for all three primers M13, 33.6 and YNZ-22 for preliminary PCR optimization step is shown in Table 3.2. The PCR reaction conditions used to generate *N. cadamba* and *N. macrophylla* DNA profiles using M13 universal primer is shown in Table 3.3.

Table 3.3: A thermal cycling profile used for preliminary PCR optimization step

Parameter	Temperature (°C)	Timing (min)	
Initial Denaturation	94.0	2	
Denaturation	94.0	1	} 35 cycles
Annealing	48.0	1	
Extension	72.0	2	
Final Extension	72.0	10	
Hold	4.0	∞	

Table 3.4: PCR reaction condition with M13 minisatellite universal primer.

Reagent	Concentration	Volume (μl)
PCR buffer	1×	2.5
dNTPs	2 mM	2.5
MgCl ₂	50 mM	1.5
Primer	2.5 pmol	6.0
<i>Taq</i> DNA polymerase	0.5 U	4.0
Template DNA	15 ng/ μl	1.0
ddH ₂ O		7.5
Total volume		25.0